

ANTI- STRETCH MARK CREAM

	COMMERCIAL NAME	INCI NAME	%
A1	COCO CAPRYLATE	COCO CAPRYLATE	6,75
A2	OLIVOIL GLUTAMATE EMULSIFIER	SODIUM OLIVOYL GLUTAMATE, CETEARYL ALCOHOL, GLYCERYL STEARATE	7,50
A3	CAPRYLIC/CAPRIC TRIGLYCERIDE	CAPRYLIC/CAPRIC TRIGLYCERIDE	5,25
A4	CETYL ALCOHOL	CETYL ALCOHOL	3,75
A5	SHEA BUTTER	BUTYROSPERMUM PARKII BUTTER	0,20
A6	TOCOPHERYL ACETATE	TOCOPHERYL ACETATE	1,75
B1	DEMINERALIZED WATER	AQUA	44,98
B2	KEMIDERM NMF V	UREA, GLUCOSE, FRUCTOSE, HYDROLYZED WHEAT PROTEIN, SODIUM GLUTAMATE, GLYCINE, LYSINE, MALIC ACID, TARTARIC ACID, CITRIC ACID, GLYCOLIC ACID, LACTIC ACID, SODIUM PCA, HYDROGENATED STARCH HYDROLYSATE, AQUA	3,00
B3	PANTHENOL	PANTHENOL	1,00
B4	GLYCERIN	GLYCERIN	2,70
B5	CHONDROITIN SULFATE	SODIUM CHONDROITIN SULFATE	0,05
B6	ALLANTOIN	ALLANTOIN	0,05
C1	KALINAT AW	SODIUM DNA	0,10
C2	DEMINERALIZED WATER	AQUA	5,00
D	CIKADERM	CROTON LECHLERI RESIN EXTRACT, ALUMINUM SUCROSE OCTASULFATE, GLYCERIN, AQUA	3,00
E	SODIUM HYALURONATE 1% SOLUTION	AQUA, SODIUM HYALURONATE	10,00
F	DOW CORNING 1501 - ABIL OSW 5	CYCLOPENTASILOXANE, DIMETHICONOL	0,50
G	ABIL B 8839 - DOW CORNING 345 FLUID	CYCLOPENTASILOXANE	1,00
H	VITAMINE A PALMITATE	RETINYL PALMITATE	0,02
I	VITAMINE F	LINSEED ACID	0,20
J1	IVY DRY EXTRACT	HEDERA HELIX LEAF EXTRACT	0,10
J2	DEMINERALIZED WATER	AQUA	1,00
K1	ASIAN CENTELLA DRY EXTRACT	CENTELLA ASIATICA LEAF EXTRACT	0,10
K2	DEMINERALIZED WATER	AQUA	1,00
L	BLACK CURRANT GLYCOLIC EXTRACT	RIBES NIGRUM FRUIT EXTRACT, PROPYLENE GLYCOL	0,50
M	PASSIONFLOWER GLYCOLIC EXTRACT	PASSIFLORA INCARNATA FLOWER EXTRACT, PROPYLENE GLYCOL	0,50
N	PRESERVATIVES, PARFUM and pH REGULATOR	PRESERVATIVES, PARFUM and pH REGULATOR	AS NEEDED
O	STABILIZER/THICKENER	STABILIZER/THICKENER	AS NEEDED

Heat all of the ingredients of phase A to a temperature of 72°C +/- 2° C mixing.

Mix the ingredients of phase B until a limpid, transparent solution is obtained. Heat phase B to a temperature of 75° +/- 2°C

When the indicated temperatures have been reached, add phase B to phase A while stirring and homogenizing.

After the union, continue to stir and homogenize for about 10 minutes. Cool the product to a temperature of 40°C, then add the phases from C to M one by one, homogenizing for a few minutes after every addition.

All the powdered ingredients (dry extracts and C1) must be dissolved into water before the addition.

Stop homogenizing and continue stirring until the emulsion reaches room temperature.

Add O as needed and set the pH between 5,90 and 6,30.

In the end, add STABILIZER/THICKENER as much as needed to get the desired viscosity.