

	COMMERCIAL NAME	INCI NAME	%
A1	OLIVOIL EMULSIFIER FOKSNaB	OLIVOYL HYDROLYZED WHEAT PROTEIN , CETEARYL ALCOHOL, GLYCERYL STEARATE, GLYCERYL OLEATE	10,00
A2	CAPRYLIC/CAPRIC TRIGLYCERIDE	CAPRYLIC/CAPRIC TRIGLYCERIDE	5,00
A3	ETHYLHEXYL PALMITATE	ETHYLHEXYL PALMITATE	6,00
A4	CETYL ALCOHOL	CETYL ALCOHOL	4,00
A5	CETEARYL ETHYLHEXANOATE	CETEARYL ETHYLHEXANOATE	3,00
A6	TOCOPHERYL ACETATE	TOCOPHERYL ACETATE	1,00
A7	DIMETHICONE 350 CPS	DIMETHICONE	1,00
A8	AVOBENZONE	BUTYL METHOXYDIBENZOYLMETHANE	0,50
A9	OMC	ETHYLHEXYL METHOXYCINNAMATE	3,00
B1	DEMINERALIZED WATER	AQUA	11,15
B2	KEMIDERM NMF V	UREA, GLUCOSE, FRUCTOSE, HYDROLYZED WHEAT PROTEIN, SODIUM GLUTAMATE, GLYCINE, LYSINE, MALIC ACID, TARTARIC ACID, CITRIC ACID, GLYCOLIC ACID, LACTIC ACID, SODIUM PCA, HYDROGENATED STARCH HYDROLYSATE, AQUA	3,00
B3	GLYCEROL	GLYCERIN	3,00
C1	EUSOLEX 232 - PARSOL HS - UVISOL PBSA	PHENYLBENZIMIDAZOLE SULFONIC ACID	2,80
C2	DEMINERALIZED WATER	AQUA	15,00
D1	SODIUM HYDROXIDE	SODIUM HYDROXIDE	0,50
D2	DEMINERALIZED WATER	AQUA	4,00
E1	RONAFLAIR BALANCE GOLD	TITANIUM DIOXIDE, MICA, TIN DIOXIDE	1,00
E2	RONAFLAIR BALANCE RED	TITANIUM DIOXIDE, MICA, TIN DIOXIDE	4,00
F	APALIGHT PF	AQUA, HYDROXYAPATITE	10,00
G1	KALILIGHT EMS	HYDROXYAPATITE, CYSTEINE, GLUTATHIONE, ZEOLITE, ETHOXYDIGLYCOL	1,00
G2	DEMINERALIZED WATER	AQUA	6,00
H1	KALINAT AW POWDER	SODIUM DNA	0,05
H2	DEMINERALIZED WATER	AQUA	2,00
I	SODIUM HYALURONATE 1% SOLUTION	AQUA, SODIUM HYALURONATE	2,50
J	DOW CORNING 1501 - ABIL OSW 5	CYCLOPENTASILOXANE, DIMETHICONOL	0,50
K	PRESERVATIVES, PARFUM and LATOR	pH REGU- PRESERVATIVES, PARFUM and pH REGULATOR	AS NEEDED
L	STABILIZER/THICKENER	STABILIZER/THICKENER	AS NEEDED

Heat all of the ingredients of phase A to a temperature of 72°C +/- 2° C mixing. Mix the ingredients of phase B until a limp, transparent solution is obtained. Disperse C1 in C2 and solubilize D1 in D2.

Add phase D to phase C little by little, mixing until a clear solution is obtained. Add this solution (C+D) to phase B. Add the pigments (E1 and E2) to the solution (B+C+D) and mix until a homogeneous dispersion is obtained. Heat the watery phase (B+C+D+E) to a temperature of 75° +/- 2°C. When the indicated temperatures have been reached, add the watery to phase to the oily phase while stirring and homogenizing. After the union, continue to stir and homogenize for about 10 minutes. Cool the product to

a temperature of 40°C, then add the other phases one by one, mixing after every addition.

G1 must be pre-dispersed in G2. H1 must be solubilized in H2.

Stop homogenizing and continue stirring until the emulsion reaches room temperature.

Add preservatives, parfum and pH regulator as much as needed to set the pH between 7,00 and 7,50.

In the end, add STABILIZER/THICKENER as much as needed to get the desired viscosity.